

## Dosing Information for Healthcare Professionals

Infants' **TYLENOL** Children's **TYLENOL**

Use this chart to determine the proper dose of **TYLENOL**® for your patient. If possible, use weight to dose; otherwise use age.

**DOSE: Every 4 hours as needed.  
Do not give more than 5 doses in 24 hours.**

WEIGHT	AGE	Infants' <b>TYLENOL</b> ® Oral Suspension Active Ingredient: Acetaminophen 160 mg (in each 5 mL)	Children's <b>TYLENOL</b> ® Oral Suspension Active Ingredient: Acetaminophen 160 mg (in each 5 mL or 1 tsp)
6-11 lbs	0-3 mos	1.25 mL	—
12-17 lbs	4-11 mos	2.5 mL	—
18-23 lbs	12-23 mos	3.75 mL	—
24-35 lbs	2-3 yrs	5 mL	5 mL (1 tsp)
36-47 lbs	4-5 yrs	—	7.5 mL (1½ tsp)
48-59 lbs	6-8 yrs	—	10 mL (2 tsp)
60-71 lbs	9-10 yrs	—	12.5 mL (2½ tsp)
72-95 lbs	11 yrs	—	15 mL (3 tsp)

mL = milliliter

tsp = teaspoonful

### Remind parents and caregivers to:

- Read and follow the label on all **TYLENOL**® products
- Take every 4 hours as needed.
- Do NOT exceed more than 5 doses in 24 hours
- Do NOT use with any other product containing acetaminophen
- Use only the dosing device that comes with a specific product

All Infants' **TYLENOL**® and Children's **TYLENOL**® Oral Suspension products have the same acetaminophen concentration (160 mg/5 mL)

## Dosing Information for Healthcare Professionals

Concentrated  
Children's **MOTRIN** Motrin  
Infants' Drops

Use this chart to determine the proper dose of **MOTRIN**® for your patient. If possible, use weight to dose; otherwise use age.

**DOSE: Every 6-8 hours as needed.  
Do not give more than 4 doses in 24 hours.**

WEIGHT	AGE	Concentrated <b>MOTRIN</b> ® Infants' Drops Active Ingredient: Ibuprofen 50 mg (NSAID) <sup>†</sup> (in each 1.25 mL)	Children's <b>MOTRIN</b> ® Oral Suspension Active Ingredient: Ibuprofen 100 mg (NSAID) <sup>†</sup> (in each 5 mL or 1 tsp)
6-11 lbs	0-5 mos	—	—
12-17 lbs	6-11 mos	1.25 mL	—
18-23 lbs	12-23 mos	1.875 mL	—
24-35 lbs	2-3 yrs	—	5 mL (1 tsp)
36-47 lbs	4-5 yrs	—	7.5 mL (1½ tsp)
48-59 lbs	6-8 yrs	—	10 mL (2 tsp)
60-71 lbs	9-10 yrs	—	12.5 mL (2½ tsp)
72-95 lbs	11 yrs	—	15 mL (3 tsp)

mL = milliliter

tsp = teaspoonful

### Remind parents and caregivers to:

- Read and follow the label on all **MOTRIN**® products
- Take every 6-8 hours as needed. Do NOT exceed more than 4 doses in 24 hours
- Do NOT administer longer than 10 days, unless directed by a doctor
- Ask a doctor or pharmacist before use if the child is taking any other drug containing an NSAID<sup>†</sup> (prescription or non-prescription)  
— **MOTRIN**® contains ibuprofen
- Use only the dosing device that comes with a specific product

<sup>†</sup>Nonsteroidal anti-inflammatory drug

©McNEIL-PPC, Inc. 2013 • CTCM-0144

## Dosing Information for Healthcare Professionals

Concentrated  
**Children's Motrin**  
Motrin  
Infants' Drops

Use this chart to determine the proper dose of MOTRIN<sup>®</sup> for your patient. If possible, use weight to dose; otherwise use age.

**DOSE: Every 6-8 hours as needed.  
Do not give more than 4 doses in 24 hours.**

WEIGHT	AGE	Concentrated MOTRIN <sup>®</sup> Infants' Drops Active Ingredient: Ibuprofen 50 mg (NSAID) <sup>1</sup> (in each 1.25 mL)	Children's MOTRIN <sup>®</sup> Oral Suspension Active Ingredient: Ibuprofen 100 mg (NSAID) <sup>1</sup> (in each 5 mL or 1 tsp)
6-11 lbs	0-5 mos	—	—
12-17 lbs	6-11 mos	1.25 mL	—
18-23 lbs	12-23 mos	1.875 mL	—
24-35 lbs	2-3 yrs	—	5 mL (1 tsp)
36-47 lbs	4-5 yrs	—	7.5 mL (1½ tsp)
48-59 lbs	6-8 yrs	—	10 mL (2 tsp)
60-71 lbs	9-10 yrs	—	12.5 mL (2½ tsp)
72-95 lbs	11 yrs	—	15 mL (3 tsp)

mL = milliliter

tsp = teaspoonful

### Remind parents and caregivers to:

- Read and follow the label on all MOTRIN<sup>®</sup> products
- Take every 6-8 hours as needed. Do NOT exceed more than 4 doses in 24 hours
- Do NOT administer longer than 10 days, unless directed by a doctor
- Ask a doctor or pharmacist before use if the child is taking any other drug containing an NSAID<sup>1</sup> (prescription or non-prescription)  
— MOTRIN<sup>®</sup> contains ibuprofen
- Use only the dosing device that comes with a specific product

<sup>1</sup>Nonsteroidal anti-inflammatory drug

©McNEIL-PPC, Inc. 2013 • CTCM-0144

## Dosing Information for Healthcare Professionals

Infants' **TYLENOL** Children's **TYLENOL**

Use this chart to determine the proper dose of TYLENOL<sup>®</sup> for your patient. If possible, use weight to dose; otherwise use age.

**DOSE: Every 4 hours as needed.  
Do not give more than 5 doses in 24 hours.**

WEIGHT	AGE	Infants' TYLENOL <sup>®</sup> Oral Suspension Active Ingredient: Acetaminophen 160 mg (in each 5 mL)	Children's TYLENOL <sup>®</sup> Oral Suspension Active Ingredient: Acetaminophen 160 mg (in each 5 mL or 1 tsp)
6-11 lbs	0-3 mos	1.25 mL	—
12-17 lbs	4-11 mos	2.5 mL	—
18-23 lbs	12-23 mos	3.75 mL	—
24-35 lbs	2-3 yrs	5 mL	5 mL (1 tsp)
36-47 lbs	4-5 yrs	—	7.5 mL (1½ tsp)
48-59 lbs	6-8 yrs	—	10 mL (2 tsp)
60-71 lbs	9-10 yrs	—	12.5 mL (2½ tsp)
72-95 lbs	11 yrs	—	15 mL (3 tsp)

mL = milliliter

tsp = teaspoonful

### Remind parents and caregivers to:

- Read and follow the label on all TYLENOL<sup>®</sup> products
- Take every 4 hours as needed.
- Do NOT exceed more than 5 doses in 24 hours
- Do NOT use with any other product containing acetaminophen
- Use only the dosing device that comes with a specific product

All Infants' TYLENOL<sup>®</sup> and Children's TYLENOL<sup>®</sup> Oral Suspension products have the same acetaminophen concentration (160 mg/5 mL)